

## Patient Pre-Inclusion Assessment Form for The Insides® System

Date of double enterostomy surgery or discovery of enterocutaneous fistula		
Description of the remaining bowel and location of the double enterostomy or enteroatmospheric fistula (proximal and distal orifice)		
<b>Inclusion criteria</b>		
1	Double enterostomy or enteroatmospheric fistula with proximal and distal outlets that can be contained in one ostomy appliance.	
2	Be able to install a feeding tube into the distal limb that has a minimum size of 20 Fr. Please speak with the clinical team for any patient that falls outside of this range	
3	Minimum length of 20 cm of the downstream segment	
4	Opacification/contrast image of the distal limb to exclude stenosis, narrowing, obstruction, or anastomotic leak	
5	Minimum ostomy output of 1000 ml/24h (while on anti-motility medication and oral food)	
6	Ostomy appliance is changed a minimum of 24 hourly (the appliance is not leaking consistently)	
7	High involvement/availability of the multidisciplinary team	
8	Adequate dentition to chew food properly or ability to blend food	
9	Retained cognitive ability of the patient or has constant carer available to manage therapy	
10	Adequate patient education so that the patient is ready to start treatment and motivated to receive the benefits of chyme reinfusion therapy	
11	Family and friend support	
<b>Exclusion until resolved</b>		
1	Fever $\geq 38^\circ$ .	
2	Sepsis	
3	Intraperitoneal collection	
4	Peritoneal carcinosis unless used in palliative treatment	
5	Impaired dentition or edentulism	
6	Impaired cognitive abilities	
7	Serious psychiatric disorders	
8	Limited availability of the multidisciplinary team	
9	Clostridium Difficile Infection	